

**A Pediatric Guide to
Children's Oral Health
Flip Chart**

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™



Primary Teeth

TEETH FACTS

- Primary teeth are also called baby teeth.
- By age 3 years, there are usually 20 primary teeth.
- The spacing between children's baby teeth is important because it allows enough room for the bigger, permanent teeth.
- Primary teeth have thinner enamel and appear whiter (translucent/almost bluish) than permanent teeth.
- Disease may progress more quickly in primary teeth.
- The biting surfaces of posterior teeth are grooved and pitted.
- Permanent teeth have wavy edges (mamelons) when they erupt, which smooth out with normal wear and tear.

TOOTH ERUPTION AND SHEDDING/EXFOLIATION

- The first tooth eruption is usually between 4 and 15 months of age.
- If eruption of the first tooth has not occurred by 18 months, the child should be referred to a dentist for evaluation.
- Premature and low birth weight babies can have delayed primary tooth eruption and enamel defects, putting them at higher risk for decay.
- Eruption is usually symmetrical (lower teeth usually before upper) in the following pattern for primary teeth: central incisors, lateral incisors, first molars, canines, second molars. Exfoliation often follows a similar pattern.
- A helpful mnemonic to remember the timing of primary eruption is the 7+4 rule. At 7 months of age, children should have their first teeth; at 11 months (4 months later), they should have 4 teeth. At 15 months of age (4 months later), they should have 8 teeth; at 19 months, they should have 12 teeth; at 23 months, they should have 16 teeth; and at 27 months, they should have 20 teeth.

- Eruption is similar for the permanent teeth, beginning between 5 and 7 years and usually finishing by 13 to 14 years of age. The typical pattern is central incisors, lateral incisors, first molars, premolars, canines, second molars, and third molars (wisdom teeth), although not everyone develops or erupts third molars.
- It is common to see permanent teeth erupt behind the primary incisor teeth in the lower jaw. This typically resolves itself without intervention, although professional dental monitoring is indicated.
- The first permanent molars erupt around 6 years of age.
- Teeth will sometimes erupt entirely out of the "normal" anticipated sequence; this should not be a concern.
- Tooth loss (also known as shedding or exfoliation) usually starts with the lower primary central incisors.

See *A Pediatric Guide to Children's Oral Health: Reference Guide* for more information about potential tooth abnormalities and development in children with special needs.

Primary Teeth



PRIMARY TEETH

		Upper Teeth	Erupt	Shed
		Central incisor	8-12 mos.	6-7 yrs.
		Lateral incisor	9-13 mos.	7-8 yrs.
		Canine (cuspid)	16-22 mos.	10-12 yrs.
		First molar	13-19 mos.	9-11 yrs.
		Second molar	25-33 mos.	10-12 yrs.
		Lower Teeth	Erupt	Shed
		Second molar	23-31 mos.	10-12 yrs.
		First molar	14-18 mos.	9-11 yrs.
		Canine (cuspid)	17-23 mos.	9-12 yrs.
		Lateral incisor	10-16 mos.	7-8 yrs.
	Central incisor	6-10 mos.	6-7 yrs.	

Early Childhood Caries

Early childhood caries (ECC) is a transmissible infectious process that affects children younger than 5 years and results in severe decay and tooth destruction. The previous terminology, *baby bottle tooth decay* and *nursing caries*, has been replaced with the term ECC because we now understand that the process of caries is independent of the route of feeding but is dependent on the frequency of refined carbohydrates in the diet. Early childhood caries is a particularly virulent form of caries that spreads rapidly within the mouth and typically results in severe dental disease.

NORMAL HEALTHY PRIMARY TEETH

- Teeth should be white with smooth surfaces.
- Spacing between teeth is desirable.
- Gum tissue (gingival mucosa) should be smooth, pink, firm, and immobile.

CHALKY WHITE SPOTS

- First appear as dull white bands on the smooth surface of the tooth at the gum line, followed by yellow or brown discoloration.
- Are indicative of early decay or weakened enamel.
- May be reversible with exposure to topical fluoride and plaque removal.
- Indicate referral to a dental home as soon as possible.
- If left untreated, will lead to cavitation.

BROWN SPOTS

- Soft brown or black spots appear on the tooth, with progression slowly toward the chewing surface of the tooth.
- Put the affected tooth at risk for fracture.
- Signify advanced or severe decay.
- Dental referral is crucial!

SEVERE DECAY

- Refers to the presence of multiple cavities in several teeth.
- May lead to early tooth loss and affects a child's ability to chew food and his or her self-esteem.
- Often requires general anesthesia in the operating room to remediate.
- Places children at risk for infection and cellulitis.

Children at High Risk for Early Tooth Decay


- Children on Medicaid
- Children whose mother or primary caregiver has cavities
- Children with siblings who have cavities
- Premature or low birth weight children
- Children with special health care needs
- Children who use a bottle after 15 months of age or have sweets and starchy snacks more than 3 times a day

Early Childhood Caries

**Normal Healthy
Primary Teeth**



**Chalky
White Spots**



Brown Spots



Severe Decay



Feeding

BREASTFEEDING

- Although breast milk alone is not cariogenic, it may become cariogenic when combined with other carbohydrate sources.
- When a child is breastfed on demand, with high frequency and duration at night, it is important to implement oral hygiene following feedings.
- Stop night feedings once teeth erupt. The majority of infants are physiologically able to tolerate a prolonged fast around 6 months of age, which is when teeth typically begin to erupt.

BOTTLE-FEEDING

- Bottles should only be used with formula, breast milk, or water. Fruit juices, soft drinks, sweet teas, formula, or milk should not be put in a baby's bottle during bedtime or nap time. At these times, bottles should only contain water.
- Infants should be held when bottle-fed. If a bottle is given with anything other than water at nap time or bedtime, parents should use a cloth to wipe the baby's mouth prior to laying the baby down.
- Bottles should not be propped with infants in cribs or car seats. Prolonged and frequent exposure to sugary liquids contributes to the caries process, and children who drink bottles while lying down may be more prone to ear infections.

SIPPY CUP

- Introduce a cup as soon as the infant can sit unsupported (around 6 months of age) and try to eliminate the bottle by 1 year of age.
- Sippy cups containing fruit juices, soft drinks, sweet teas, formula, or milk should not be given to the child at bedtime or nap time.
- If the sippy cup is offered between meals, it should contain only milk or water. If juice is offered, it should be restricted to mealtimes.
- Other sugar-containing drinks should be avoided.

FOOD GUIDANCE

- Provide the child with healthy alternatives such as fruits and vegetables cut into small pieces (to avoid choking) or whole grain snacks.
- Pre-tasting, pre-chewing, and sharing of utensils should be avoided because bacteria is transmitted through saliva.
- Avoid sticky foods like raisins, fruit leather, and hard candies.
- Discourage grazing.

Feeding

Breastfeeding




Bottle-feeding



Sippy Cup



Food Guidance



Nonnutritive Sucking

Sucking is a normal baby reflex. It helps babies feel secure and happy and helps them learn about their world. Babies may suck their thumbs, fingers, or hands, or a pacifier or other inanimate object such as a blanket or toy. Most children discontinue their nonnutritive sucking habit between the ages of 2 and 4 years.

THUMB AND FINGER SUCKING

- Babies who suck their thumbs usually continue the habit longer than pacifier users.
- Prolonged thumb sucking may cause problems with proper growth of the mouth and the alignment of teeth. It also can cause changes in the roof of the mouth.
- The most common dental effect of nonnutritive sucking is anterior, upward movement of the maxillary central incisors and palatal bone, which may result in an anterior open bite. Other possible effects include maxillary constriction and posterior crossbite.
- Children should be encouraged to discontinue their nonnutritive sucking habits by 4 years of age.

PACIFIER USE

- Pacifiers should never be dipped in sugary substances such as honey and sugar.
- Pacifier use during sleep is associated with a decreased incidence of sudden infant death syndrome.
- Pacifiers should never be used to replace or delay meals and should be offered only when the caregiver is certain the child is not hungry.
- Pacifiers should have ventilation holes and a shield wider than the child's mouth (at least 1 ¼ inches in diameter).
- Pacifiers should be one piece and made of a durable material, replaced when worn, and never tied by a string to the crib or around a child's neck or hand.
- Physiologic pacifiers are preferable to conventional pacifiers because they may have less dental effects.

TEETHING

- Teething is the emergence of the first primary (baby) teeth through a baby's gums.
- Teething begins as early as 3 months and continues until the child is approximately 3 years of age.
- Some babies may have tender and swollen gums when teething.
- Diarrhea, rashes, and a fever are not normal for a teething baby.

INSTRUCTIONS TO PARENTS

- Remove the drool on the baby's face to prevent rashes from developing.
- Give the baby something hard or cold to chew on, making sure it is big enough that it can't be swallowed or break into small pieces. Examples include refrigerated teething rings, pacifiers, spoons, clean wet washcloths, and frozen bagels or bananas.
- Gently rub the baby's gums with a clean finger.
- If the baby seems irritable, acetaminophen can be used.
- Topical teething gels sold over the counter (OTC) are sometimes used for teething. However, these gels can carry serious risks, including local reactions, seizures (with overdose), and methemoglobinemia. Parents should be instructed on proper dosing of OTC analgesic medications.
- Regularly disinfect teething rings and objects and wash hands to avoid gastrointestinal disturbances and infections.

Nonnutritive Sucking

Thumb Sucking




Finger Sucking



Pacifier Use



Teething



Fluoride Modalities

Fluoride is a naturally occurring substance that is safe and effective in preventing tooth decay.

FLUORIDATED WATER

- It can be found in communities that supplement tap water with fluoride.
- The Centers for Disease Control and Prevention (CDC) My Water's Fluoride Web site (<http://apps.nccd.cdc.gov/MWF/Index.asp>) allows consumers in currently participating states to learn the fluoridation status of their water system.
- Fluoride can be found in some bottled water. Look for fluoride content on the label.

FLUORIDE SUPPLEMENTS (DROPS AND TABLETS)

American Dental Association Dietary Fluoride Supplement Schedule for Children at High Caries Risk

(http://ebd.ada.org/contentdocs/6327_Fluoride_Chairside_Tool.pdf)

Age	<0.3 ppm	0.3–0.6 ppm	>0.6 ppm
0–6 mo	None	None	None
6 mo–3 y	0.25 mg/d	None	None
3–6 y	0.50 mg/d	0.25 mg/d	None
6–16 y	1.0 mg/d	0.50 mg/d	None

- Supplements can be used as early as 6 months of age if a child does not have access to an adequate amount of fluoride in the community water supply.
- For children at low caries risk, supplements are not recommended. Other sources of fluoride should be considered to prevent caries.
- For children at high caries risk, supplements are recommended according to the American Dental Association's schedule.
- When fluoride supplements are prescribed they should be taken daily to maximize benefit.
- If a family uses well water, be sure to order well water fluoride testing before prescribing supplements.
- Specific guidelines are used to determine the amount of fluoride needed based on a child's age.
- Drops come in 0.5 mg/mL.
- The topical effect of tablets is the preferred route. Many children can be transitioned to chew tablets as early as 15 to 18 months of age.

FLUORIDATED TOOTHPASTE

- A smear can be used as soon the first tooth erupts if the child is at high risk for caries.
- Caution is advised when using fluoridated toothpaste for young children because they may swallow excessive amounts of toothpaste.
- Unless the child has less than optimal exposure to fluoride, fluoridated toothpaste should not be used until the child is 2 years of age. A pea-sized amount of fluoride-containing toothpaste (shown on opposite side in photo of a child's toothbrush) has the recommended total daily intake of fluoride for a 2-year-old child.
- Children younger than 6 years should use only a pea-sized amount of fluoridated toothpaste.
- Check the fluoride content of toothpastes; almost all toothpastes manufactured in the United States provide topical fluoride, but not all natural toothpastes do.

FLUORIDE VARNISH

- Most fluoride varnishes are lacquers containing 5% sodium fluoride in a colophony/resin base.
- Fluoride varnish is painted on a child's teeth by a dentist or child health care professional if the child is at high risk for caries.
- The varnish protects teeth from decay and can arrest the progression of a chalky white spot lesion to a full-blown cavity.

Threat of Fluorosis

- Fluorosis is a condition caused by an excessive intake of fluoride.
- The effects of fluorosis are mainly aesthetic. Mild fluorosis results in lacy markings on the tooth's enamel surface; in moderate fluorosis, a white opacity is easy to see over 50% of the tooth; and severe fluorosis results in pitted, brittle enamel.
- Threat of fluorosis disappears after mineralization of all permanent teeth (usually 8 years of age).



Fluoride Modalities

Fluoridated Water



Fluoride Supplements



Fluoridated Toothpaste



Baby toothbrush with a smear of toothpaste



Child's toothbrush with a pea-sized amount of toothpaste

Fluoride Varnish



Toothbrushing

FLUORIDE TOOTHBRUSHING RECOMMENDATIONS

Age	Toothbrushing Recommendations
<1 y	Clean teeth with soft toothbrush.
1–2 y	Parent should perform brushing.
2–6 y	Pea-sized amount of fluoride-containing toothpaste 2 times per day; parent performs or supervises.
>6 y	Brush with fluoridated toothpaste 2 times per day.

ADVICE TO PARENTS

- Clean or brush a young child's teeth twice daily.
- Begin wiping the gums of even a very small infant with a soft washcloth or soft toothbrush, even prior to tooth eruption, to establish a daily oral hygiene routine.
- Toothbrushes for infants and toddlers should be soft with a small head and a large handle.
- Toothbrushing should be supervised until the child can reliably rinse and spit out excess toothpaste (usually 6 years of age). Younger children do not have the hand coordination necessary for independent toothbrushing prior to that age.
- Electric toothbrushes are especially useful in situations of limited movement. They do the work for you, they position well, and the small head can help limit the amount of toothpaste to what is appropriate for children.
- All accessible surfaces of each tooth need to be brushed.
- Remind parents to not allow their child to swallow fluoridated toothpaste.

FLOSSING RECOMMENDATIONS

Flossing is an essential part of the tooth-cleaning process. It removes food particles and plaque between teeth that brushing misses.


ADVICE TO PARENTS

- Flossing should begin when 2 teeth touch, typically between 2 and 2½ years of age. Some children may only need a few back teeth flossed and others may need flossing between all their tight teeth, depending on dental spacing.
- Children usually need assistance with flossing until they are 8 to 10 years of age.
- Flossing tools, such as pre-threaded flossers or floss holders, may be helpful for children who are just learning how to floss.
- Some children may find it easier to use a loop of floss, which is created by taking a piece of floss about 10 inches long and tying the ends together into a circle. Parents (and older children) can hold the floss tightly between the thumbs and forefingers to floss.

Toothbrushing



Flossing



Fluoride Varnish

FLUORIDE VARNISH FACTS

- Concentrated topical fluoride.
- Five-percent sodium fluoride in a resin base.
- Effective in preventing early childhood caries and reducing caries progression.
- Fluoride application by physicians may be a billable procedure. Check with the local Medicaid program for more information.

APPLYING FLUORIDE VARNISH

- Have everything ready.
- Open varnish packet and mix it well.
- Wipe child's teeth dry with a clean gauze.
- Paint child's teeth with varnish using disposable applicator.
- Instruct parents.
 - ♦ *Do not brush the child's teeth until the next day.*
 - ♦ *The child's teeth may be slightly yellow until they are brushed.*
 - ♦ *The child can eat and drink right away but should avoid hot liquids.*



FREQUENCY OF APPLICATION

- For moderate caries risk, apply every 6 months.
- For high caries risk, children may benefit from increased application frequency of every 3 to 4 months.
- Payment codes may only be allowed for 2 to 3 applications per year. (Check with the local Medicaid program for information about payment.)

CHILDREN AGED 0 TO 3 YEARS WHO WILL BENEFIT FROM FLUORIDE VARNISH

- All children eligible for Medicaid
- Siblings with cavities before 6 years of age
- Premature children
- Children with special health care needs
- Children who use a bottle after 15 months or have sweet or starchy snacks more than 3 times a day
- Children without a dental home

Fluoride Varnish



ACKNOWLEDGMENTS

A Pediatric Guide to Children's Oral Health is a collaborative project of the Oral Health Initiative coordinated by the American Academy of Pediatrics (AAP) and the federal Maternal and Child Health Bureau.

We request the following citation be used when referencing this document:

American Academy of Pediatrics. *A Pediatric Guide to Children's Oral Health*. Elk Grove Village, IL: American Academy of Pediatrics; 2009

For information about the AAP Oral Health Initiative, contact

American Academy of Pediatrics
141 Northwest Point Blvd
Elk Grove Village, IL 60007-1098
E-mail: oralhealth@aap.org
Web site: www.aap.org/oralhealth

The recommendations in this publication do not indicate an exclusive course of treatment or serve as a standard of medical care. Variations, taking into account individual circumstances, may be appropriate.

Products are shown or mentioned for informational purposes only and do not imply an endorsement by the AAP. The AAP does not recommend any specific brand of products or services.

Copyright © 2010 American Academy of Pediatrics.
All rights reserved.



PRIMARY AUTHORS

- Rama Oskouian, DMD
- Giusy Romano-Clarke, MD, FAAP

ADDITIONAL CONTRIBUTORS

- Lauren Barone, MPH
- Suzanne Boulter, MD, FAAP
- Melinda B. Clark, MD, FAAP
- David M. Krol, MD, MPH, FAAP
- Wendy Nelson, ACCE
- Diona L. Reeves
- Rebecca Slayton, DDS, PhD

THE FOLLOWING INDIVIDUALS AND ORGANIZATIONS PROVIDED PHOTOGRAPHS FOR THIS RESOURCE:

- American Dental Association
- ANZ Photography
- Suzanne Boulter, MD, FAAP
- Melinda B. Clark, MD, FAAP
- Content Visionary
- Yasmi Crystal, DMD
- Joanna Douglas, BDS, DDS
- Donald Greiner, DDS, MSc
- Martha Ann Keels, DDS, PhD
- Rama Oskouian, DMD
- Giusy Romano-Clarke, MD, FAAP
- Tegwyn H. Brickhouse, DDS, PhD

A special thanks to AAP staff and advisory committee members who helped in the production of this resource.

This project is supported in part by Grant No. U93 MC 00184 from the Maternal and Child Health Bureau (Title V, Social Security Act), Health Resources and Services Administration, Department of Health and Human Services and the American Dental Association Foundation.



American Academy of Pediatrics
DEDICATED TO THE HEALTH OF ALL CHILDREN™

